



Amgen Announces Positive Top-Line Results From Phase 3 GAUSS-3 Trial Of Repatha® (Evolocumab) In Statin-Intolerant Patients With High Cholesterol

February 4, 2016

Study Meets Co-Primary Endpoints of LDL Cholesterol Reduction

THOUSAND OAKS, Calif., Feb. 4, 2016 /PRNewswire/ -- Amgen (NASDAQ:AMGN) today announced that the Phase 3 GAUSS-3 (Goal Achievement After Utilizing an Anti-PCSK9 Antibody in Statin Intolerant Subjects-3) trial evaluating Repatha® (evolocumab) in patients with high cholesterol who cannot tolerate statins met its co-primary endpoints: mean percent reductions from baseline in low-density lipoprotein cholesterol (LDL-C) at weeks 22 and 24, and the percent reduction from baseline in LDL-C at week 24. The mean percent reductions in LDL-C, or "bad" cholesterol, compared to ezetimibe, were consistent with results observed in the 12-week Phase 2 GAUSS-1 and Phase 3 GAUSS-2 trials.

"The positive results from the GAUSS-3 study contribute to the growing body of evidence supporting Repatha as an innovative treatment option for patients who have not been able to adequately lower their LDL cholesterol through diet and statins alone," said Sean E. Harper, M.D., executive vice president of Research and Development at Amgen. "Many patients with high LDL cholesterol are unable to tolerate effective doses of statins, and the findings from the rigorously-designed GAUSS-3 study confirm the results in the previous GAUSS studies. We look forward to exploring these data further."

GAUSS-3 is a three-part trial that is evaluating the safety, tolerability and efficacy of Repatha, an injectable proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor, in patients with high cholesterol who could not tolerate statins due to muscle-related side effects (MRSE). The active-controlled part of the trial evaluated the effect of 24 weeks of treatment with Repatha compared to ezetimibe on percent change from baseline in LDL-C.

In the GAUSS-3 trial there were no new safety findings. The most common adverse events that occurred in greater than 5 percent of patients in the Repatha group were myalgia, nasopharyngitis, muscle spasms, arthralgia, pain in extremity, fatigue, headache and back pain.

The full data results from the Phase 3 GAUSS-3 trial will be submitted to a future medical conference and for publication.

About Repatha® (evolocumab)

Repatha® (evolocumab) is a human monoclonal antibody that inhibits proprotein convertase subtilisin/kexin type 9 (PCSK9).¹ Repatha binds to PCSK9 and inhibits circulating PCSK9 from binding to the low-density lipoprotein (LDL) receptor (LDLR), preventing PCSK9-mediated LDLR degradation and permitting LDLR to recycle back to the liver cell surface. By inhibiting the binding of PCSK9 to LDLR, Repatha increases the number of LDLRs available to clear LDL from the blood, thereby lowering LDL-C levels.¹

GLAGOV, the intravascular ultrasound study, is underway to determine the effect of Repatha on coronary atherosclerosis in approximately 950 patients undergoing cardiac catheterization to test the hypothesis of robust LDL-C reduction leading to a reduction or a change in the build-up of plaque in the arteries. Results from the GLAGOV study are expected in the second half of 2016.

The FOURIER outcomes trial is designed to evaluate whether treatment with Repatha in combination with statin therapy, compared to placebo plus statin therapy, reduces the risk of cardiovascular events in patients with high cholesterol and clinically evident cardiovascular disease, and completed patient enrollment in June 2015. Top-line results from the approximately 27,500-patient event-driven FOURIER study are anticipated in the second half of 2016.

Repatha is approved in the United States, Japan, Canada, Australia, Kuwait, and in all 28 countries that are members of the European Union as well as in Norway, Iceland and Liechtenstein. Applications in other countries are pending.

Important U.S. Product Information

Repatha® is indicated as an adjunct to diet and:

- Maximally tolerated statin therapy for treatment of adults with heterozygous familial hypercholesterolemia (HeFH) or clinical atherosclerotic cardiovascular disease (ASCVD), who require additional lowering of low-density lipoprotein cholesterol (LDL-C)
- Other LDL-lowering therapies (e.g., statins, ezetimibe, LDL apheresis) in patients with homozygous familial hypercholesterolemia (HoFH) who require additional lowering of LDL-C

The effect of Repatha® on cardiovascular morbidity and mortality has not been determined.

The safety and effectiveness of Repatha® have not been established in pediatric patients with HoFH who are younger than 13 years old.

The safety and effectiveness of Repatha® have not been established in pediatric patients with primary hyperlipidemia or HeFH.

Important Safety Information

Contraindication: Repatha® is contraindicated in patients with a history of a serious hypersensitivity reaction to Repatha®.

Allergic reactions: Hypersensitivity reactions (e.g. rash, urticaria) have been reported in patients treated with Repatha®, including some that led to discontinuation of therapy. If signs or symptoms of serious allergic reactions occur, discontinue treatment with Repatha®, treat according to the standard of care, and monitor until signs and symptoms resolve.

Adverse reactions: The most common adverse reactions (>5% of Repatha®-treated patients and more common than placebo) were: nasopharyngitis, upper respiratory tract infection, influenza, back pain, and injection site reactions.

In a 52-week trial, adverse reactions led to discontinuation of treatment in 2.2% of Repatha®-treated patients and 1% of placebo-treated patients. The most common adverse reaction that led to Repatha® treatment discontinuation and occurred at a rate greater than placebo was myalgia (0.3% versus 0% for Repatha® and placebo, respectively).

Adverse reactions from a pool of the 52-week trial and seven 12-week trials:

Local injection site reactions occurred in 3.2% and 3.0% of Repatha®-treated and placebo-treated patients, respectively. The most common injection site reactions were erythema, pain, and bruising. The proportions of patients who discontinued treatment due to local injection site reactions in Repatha®-treated patients and placebo-treated patients were 0.1% and 0%, respectively.

Allergic reactions occurred in 5.1% and 4.7% of Repatha®-treated and placebo-treated patients, respectively. The most common allergic reactions were rash (1.0% versus 0.5% for Repatha® and placebo, respectively), eczema (0.4% versus 0.2%), erythema (0.4% versus 0.2%), and urticaria (0.4% versus 0.1%).

Neurocognitive events were reported in less than or equal to 0.2% in Repatha®-treated and placebo-treated patients.

In a pool of placebo- and active-controlled trials, as well as open-label extension studies that followed them, a total of 1,988 patients treated with Repatha® had at least one LDL-C value <25 mg/dL. Changes to background lipid-altering therapy were not made in response to low LDL-C values, and Repatha® dosing was not modified or interrupted on this basis. Although adverse consequences of very low LDL-C were not identified in these trials, the long-term effects of very low levels of LDL-C induced by Repatha® are unknown.

Musculoskeletal adverse reactions were reported in 14.3% of Repatha®-treated patients and 12.8% of placebo-treated patients. The most common adverse reactions that occurred at a rate greater than placebo were back pain (3.2% versus 2.9% for Repatha® and placebo, respectively), arthralgia (2.3% versus 2.2%), and myalgia (2.0% versus 1.8%).

Homozygous Familial Hypercholesterolemia (HoFH): In 49 patients with homozygous familial hypercholesterolemia studied in a 12-week, double-blind, randomized, placebo-controlled trial, 33 patients received 420 mg of Repatha® subcutaneously once monthly. The adverse reactions that occurred in at least 2 (6.1%) Repatha®-treated patients and more frequently than in placebo-treated patients, included upper respiratory tract infection (9.1% versus 6.3%), influenza (9.1% versus 0%), gastroenteritis (6.1% versus 0%), and nasopharyngitis (6.1% versus 0%).

Immunogenicity: Repatha® is a human monoclonal antibody. As with all therapeutic proteins, there is a potential for immunogenicity with Repatha®.

Please contact Amgen Medinfo at 800-77-AMGEN (800-772-6436) or 844-REPATHA (844-737-2842) regarding Repatha® availability or find more information, including full Prescribing Information, at www.amgen.com and www.Repatha.com.

About Amgen Cardiovascular

Building on more than three decades of experience in developing biotechnology medicines for patients with serious illnesses, Amgen is dedicated to addressing important scientific questions to advance care and improve the lives of patients with cardiovascular disease, the leading cause of morbidity and mortality worldwide.² Amgen's research into cardiovascular disease, and potential treatment options, is part of a growing competency at Amgen that utilizes human genetics to identify and validate certain drug targets. Through its own research and development efforts, as well as partnerships, Amgen is building a robust cardiovascular portfolio consisting of several approved and investigational molecules in an effort to address a number of today's important unmet patient needs, such as high cholesterol and heart failure.

About Amgen

Amgen is committed to unlocking the potential of biology for patients suffering from serious illnesses by discovering, developing, manufacturing and delivering innovative human therapeutics. This approach begins by using tools like advanced human genetics to unravel the complexities of disease and understand the fundamentals of human biology.

Amgen focuses on areas of high unmet medical need and leverages its biologics manufacturing expertise to strive for solutions that improve health outcomes and dramatically improve people's lives. A biotechnology pioneer since 1980, Amgen has grown to be one of the world's leading independent biotechnology companies, has reached millions of patients around the world and is developing a pipeline of medicines with breakaway potential.

For more information, visit www.amgen.com and follow us on www.twitter.com/amgen.

Amgen Forward-Looking Statements

This news release contains forward-looking statements that are based on management's current expectations and beliefs and are subject to a number of risks, uncertainties and assumptions that could cause actual results to differ materially from those described. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including estimates of revenues, operating margins, capital expenditures, cash, other financial metrics, expected legal, arbitration, political, regulatory or clinical results or practices, customer and prescriber patterns or practices, reimbursement activities and outcomes and other such estimates and results. Forward-looking statements involve significant risks and uncertainties, including those discussed below and more fully described in the Securities and Exchange Commission (SEC) reports filed by Amgen, including Amgen's most recent annual report on Form 10-K and any subsequent periodic reports on Form 10-Q and Form 8-K. Please refer to Amgen's most recent Forms 10-K, 10-Q and 8-K for additional information on the uncertainties and risk factors related to our business. Unless otherwise noted, Amgen is providing this information as of Feb. 4, 2016, and expressly disclaims any duty to update information contained in this news release.

No forward-looking statement can be guaranteed and actual results may differ materially from those we project. Discovery or identification of new product candidates or development of new indications for existing products cannot be guaranteed and movement from concept to product is uncertain; consequently, there can be no guarantee that any particular product candidate or development of a new indication for an existing product will be

successful and become a commercial product. Further, preclinical results do not guarantee safe and effective performance of product candidates in humans. The complexity of the human body cannot be perfectly, or sometimes, even adequately modeled by computer or cell culture systems or animal models. The length of time that it takes for us to complete clinical trials and obtain regulatory approval for product marketing has in the past varied and we expect similar variability in the future. We develop product candidates internally and through licensing collaborations, partnerships and joint ventures. Product candidates that are derived from relationships may be subject to disputes between the parties or may prove to be not as effective or as safe as we may have believed at the time of entering into such relationship. Also, we or others could identify safety, side effects or manufacturing problems with our products after they are on the market. Our business may be impacted by government investigations, litigation and products liability claims. If we fail to meet the compliance obligations in the corporate integrity agreement between us and the U.S. government, we could become subject to significant sanctions. We depend on third parties for a significant portion of our manufacturing capacity for the supply of certain of our current and future products and limits on supply may constrain sales of certain of our current products and product candidate development.

In addition, sales of our products are affected by the reimbursement policies imposed by third-party payors, including governments, private insurance plans and managed care providers and may be affected by regulatory, clinical and guideline developments and domestic and international trends toward managed care and healthcare cost containment as well as U.S. legislation affecting pharmaceutical pricing and reimbursement. Government and others' regulations and reimbursement policies may affect the development, usage and pricing of our products. In addition, we compete with other companies with respect to some of our marketed products as well as for the discovery and development of new products. We believe that some of our newer products, product candidates or new indications for existing products, may face competition when and as they are approved and marketed. Our products may compete against products that have lower prices, established reimbursement, superior performance, are easier to administer, or that are otherwise competitive with our products. In addition, while we routinely obtain patents for our products and technology, the protection offered by our patents and patent applications may be challenged, invalidated or circumvented by our competitors and there can be no guarantee of our ability to obtain or maintain patent protection for our products or product candidates. We cannot guarantee that we will be able to produce commercially successful products or maintain the commercial success of our existing products. Our stock price may be affected by actual or perceived market opportunity, competitive position, and success or failure of our products or product candidates. Further, the discovery of significant problems with a product similar to one of our products that implicate an entire class of products could have a material adverse effect on sales of the affected products and on our business and results of operations. Our efforts to integrate the operations of companies we have acquired may not be successful. We may experience difficulties, delays or unexpected costs and not achieve anticipated benefits and savings from our ongoing restructuring plan. Our business performance could affect or limit the ability of our Board of Directors to declare a dividend or their ability to pay a dividend or repurchase our common stock.

The scientific information discussed in this news release relating to new indications is preliminary and investigative and is not part of the labeling approved by the U.S. Food and Drug Administration or European Commission for the products. The products are not approved for the investigational use(s) discussed in this news release, and no conclusions can or should be drawn regarding the safety or effectiveness of the products for these uses.

CONTACT: Amgen, Thousand Oaks
Kristen Davis: 805-447-3008 (media)
Kristen Neese: 805-313-8267 (media)
Arvind Sood: 805-447-1060 (investors)

References

1. Amgen Data on File, Investigator Brochure.
2. World Health Organization. Cardiovascular diseases (CVDs) fact sheet. <http://www.who.int/mediacentre/factsheets/fs317/en/>. Accessed January 2016.



Logo - <http://photos.pnewswire.com/prnh/20081015/AMGENLOGO>

To view the original version on PR Newswire, visit: <http://www.pnewswire.com/news-releases/amgen-announces-positive-top-line-results-from-phase-3-gauss-3-trial-of-repatha-evolocumab-in-statin-intolerant-patients-with-high-cholesterol-300215615.html>

SOURCE Amgen