



European Commission Approves BLINCYTO® (blinatumomab) In Patients With Philadelphia Chromosome Negative Minimal Residual Disease-Positive B-cell Precursor Acute Lymphoblastic Leukemia

January 22, 2019

BLINCYTO is the First and Only Therapy for Minimal Residual Disease Approved in the European Union Approval Based on Data From the Phase 2 BLAST Study, the Largest Prospective Trial in Minimal Residual Disease-Positive Acute Lymphoblastic Leukemia

THOUSAND OAKS, Calif., Jan. 22, 2019 /PRNewswire/ -- Amgen (NASDAQ:AMGN) today announced that the European Commission (EC) has approved an expanded indication for BLINCYTO® (blinatumomab) monotherapy to include adult patients with Philadelphia chromosome negative (Ph-) CD19 positive B-cell precursor acute lymphoblastic leukemia (ALL) in first or second complete remission with minimal residual disease (MRD) greater than or equal to 0.1 percent. The approval was based on data from the Phase 2 BLAST study in frontline and relapsed/refractory ALL, the largest prospective trial for MRD-positive ALL ever conducted. BLINCYTO, a bispecific CD19-directed CD3 T cell engager (BiTE®), is the first BiTE® immunotherapy to receive regulatory approval globally.

MRD refers to the presence of cancer cells that remain detectable, despite a patient having achieved complete remission by conventional assessment.¹ MRD is only measurable through the use of highly sensitive testing methods that detect cancer cells in the bone marrow with a sensitivity of at least one cancer cell in 10,000 cells – versus about one in 20 with a conventional microscope-based evaluation.¹⁻³

"This approval represents a paradigm shift in the management of ALL in the European Union, making BLINCYTO the first and only treatment with marketing authorization to include the presence of MRD," said David M. Reese, M.D., executive vice president of Research and Development at Amgen. "We are pleased that the European Commission has seen the value BLINCYTO can bring to people living with ALL and are proud to continue to deliver on our commitment to the pursuit of breakthroughs that can transform the lives of cancer patients."

The EC approval is based on data from the Phase 2 BLAST study, which found that BLINCYTO induced a complete MRD response, or no detectable MRD, in 78 percent of patients within one treatment cycle. Safety results among MRD-positive patients were consistent with the known safety profile of BLINCYTO in relapsed or refractory B-cell precursor ALL.

"Survival rates for patients that achieve MRD-negativity are significantly higher than those for patients that remain MRD-positive, underscoring the critical importance of early testing for and eliminating residual disease in patients with ALL," said Nicola Gökbüget, M.D., principal investigator for the BLAST study and head of the German Multicenter Study Group for Adult ALL located in Frankfurt, Germany. "Data from the BLAST study demonstrated that BLINCYTO is effective in eliminating detectable residual disease and showed the potential to improve relapse-free survival in this patient population. Today's approval provides physicians across Europe with a much-needed treatment option that can potentially help prevent a relapse, which is associated with a very unfortunate prognosis."

Approval via the centralized procedure allows for obtaining a marketing authorization from the EC, which is valid in all European Union (EU) and European Economic Area (EEA)-European Free Trade Association (EFTA) states (Norway, Iceland and Liechtenstein).

In March 2018, the U.S. Food and Drug Administration (FDA) approved BLINCYTO for the treatment of adults and children with B-cell precursor ALL in first or second complete remission with MRD greater than or equal to 0.1 percent.

BLINCYTO is the first immunotherapy from Amgen's BiTE® platform, an innovative approach that helps the body's immune system target cancer cells. BiTE® antibody construct technology, pioneered by Amgen, is an innovative treatment approach that helps the body's immune system attack cancer cells without the removal of immune cells from the patient. Amgen is studying a number of "off-the-shelf" investigational BiTE® immunotherapies, with distinct targets, across a range of hematologic and solid tumors.

About the BLAST Study

The BLAST study is the largest ever prospective trial in patients with MRD-positive ALL. It is an open-label, multicenter, single-arm, Phase 2 study evaluating the efficacy, safety and tolerability of BLINCYTO in adult patients with MRD-positive B-cell precursor ALL in complete hematologic remission after three or more cycles of intensive chemotherapy. Patients received continuous IV infusion of BLINCYTO 15 µg/m²/d for four weeks, followed by two weeks off. Patients received up to four cycles of treatment and could undergo hematopoietic stem cell transplantation at any time after the first cycle, if eligible. Efficacy was based on achievement of undetectable MRD within one cycle of BLINCYTO treatment and hematological relapse-free survival (RFS). Additional secondary endpoints included incidence and severity of adverse events, overall survival, time to hematological remission and duration of complete MRD response.

Results from the BLAST study were presented at the 57th American Society of Hematology (ASH) Annual Meeting & Exposition in 2015 and published in *Blood* in 2018. Long-term overall survival (OS) data results from the BLAST study were also featured in an oral presentation during the ASH 2018 Annual Meeting & Exposition on Dec. 3, 2018.

About ALL and MRD

ALL is a rapidly progressing cancer of the blood and bone marrow that occurs in both adults and children.^{4,5} Poor outcomes have been observed in patients who achieve first or second complete hematologic remission but have persistent MRD, which remains detectable at the molecular level after treatment.^{1,6} For more information about MRD, please visit AmgenOncology.com.

About BiTE® Technology

Bispecific T cell engager (BiTE®) antibody construct is an innovative technology that can be engineered to target any tumor antigen expressed by any

type of cancer. The protein molecules are designed to kill malignant cells using the patient's own immune system by bridging T cells to tumor cells. The BiTE[®] antibody construct helps connect the T cells to the targeted cell, with the intent of causing T cells to inject toxins which trigger cancer cell death (apoptosis). Amgen is developing BiTE[®] antibody constructs to uniquely (or specifically) target numerous hematologic malignancies and solid tumors.

About BLINCYTO[®] (blinatumomab)

BLINCYTO is a bispecific CD19-directed CD3 T cell engager (BiTE[®]) immunotherapy that binds to CD19 expressed on the surface of cells of B-lineage origin and CD3 expressed on the surface of effector T cells. BLINCYTO was granted breakthrough therapy and priority review designations by the FDA in 2014, and carries full approval in the U.S. for the treatment of relapsed or refractory B-cell precursor ALL in adults and children. In the U.S., BLINCYTO is also approved for the treatment of adults and children with B-cell precursor ALL in first or second complete remission with MRD greater than or equal to 0.1 percent. This indication is approved under accelerated approval based on MRD response rate and hematological relapse-free survival. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials. In the EU, BLINCYTO is indicated for the treatment of adults with Ph- relapsed or refractory B-precursor ALL and for the treatment of Ph- CD19 positive B-cell precursor ALL in first or second complete remission with MRD greater than or equal to 0.1 percent.

BLINCYTO is now approved in 57 countries, including all member countries in the EU and EEA, Canada, Japan and Australia.

Important EU BLINCYTO[®] (blinatumomab) Safety Information

This product is subject to additional monitoring in the EU. All suspected adverse reactions should be reported in accordance with the national reporting system.

The adverse reactions described in this section were identified in clinical studies of patients with B-precursor ALL (N = 843). The most serious adverse reactions that may occur during blinatumomab treatment include: infections (24.8%), neurologic events (13.8%), neutropenia/febrile neutropenia (10.1%), cytokine release syndrome (3.3%), and tumour lysis syndrome (0.7%). The most common adverse reactions were: pyrexia (69.2%), infusion-related reactions (43.4%), infections – pathogen unspecified (42.1%), headache (32.9%), anaemia (22.8%), thrombocytopenia (20.9%), febrile neutropenia (20.2%), oedema (20.0%), neutropenia (19.7%), rash (16.7%), increased liver hepatic enzymes (16.1%), bacterial infectious disorders (15.4%), tremor (15.2%), cough (15.1%), leukopenia (13.4%), back pain (13.3%), chills (13.0%), hypotension (12.8%), viral infectious disorders (12.7%), decreased immunoglobulins (12.5%), cytokine release syndrome (11.6%), tachycardia (11.3%), insomnia (10.7%), fungal infectious disorders (10.6%) and pain in extremity (10.2%).

Please refer to the Summary of Product Characteristics for full European prescribing information.

Important Safety Information Regarding BLINCYTO[®] (blinatumomab) U.S. Indication

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGICAL TOXICITIES

- **Cytokine Release Syndrome (CRS), which may be life-threatening or fatal, occurred in patients receiving BLINCYTO[®]. Interrupt or discontinue BLINCYTO[®] as recommended.**
- **Neurological toxicities, which may be severe, life-threatening or fatal, occurred in patients receiving BLINCYTO[®]. Interrupt or discontinue BLINCYTO[®] as recommended.**

Contraindications

BLINCYTO[®] is contraindicated in patients with a known hypersensitivity to blinatumomab or to any component of the product formulation.

Warnings and Precautions

- **Cytokine Release Syndrome (CRS):** CRS, which may be life-threatening or fatal, occurred in patients receiving BLINCYTO[®]. The median time to onset of CRS is 2 days after the start of infusion. Closely monitor patients for signs and symptoms of serious adverse events such as fever, headache, nausea, asthenia, hypotension, increased alanine aminotransferase (ALT), increased aspartate aminotransferase (AST), increased total bilirubin (TBILI), and disseminated intravascular coagulation (DIC). The manifestations of CRS after treatment with BLINCYTO overlap with those of infusion reactions, capillary leak syndrome (CLS), and hemophagocytic histiocytosis/macrophage activation syndrome (MAS). In clinical trials of BLINCYTO, CRS was reported in 15% of patients with relapsed or refractory ALL and in 7% of patients with MRD-positive ALL. Interrupt or discontinue BLINCYTO[®] for evidence of CRS, as outlined in the PI.
- **Neurological Toxicities:** Approximately 65% of patients receiving BLINCYTO[®] in clinical trials experienced neurological toxicities. The median time to the first event was within the first 2 weeks of BLINCYTO[®] treatment, and the majority of events resolved. The most common ($\geq 10\%$) manifestations of neurological toxicity were headache and tremor. Severe, life-threatening, or fatal neurological toxicities occurred in approximately 13% of patients, including encephalopathy, convulsions, speech disorders, disturbances in consciousness, confusion and disorientation, and coordination and balance disorders. Manifestations of neurological toxicity included cranial nerve disorders. Monitor patients for signs or symptoms of neurological toxicity and interrupt or discontinue BLINCYTO[®] as outlined in the PI.
- **Infections:** Approximately 25% of patients receiving BLINCYTO[®] in clinical trials experienced serious infections such as sepsis, pneumonia, bacteremia, opportunistic infections, and catheter-site infections, some of which were life-threatening or fatal. Administer prophylactic antibiotics and employ surveillance testing as appropriate during treatment. Monitor patients for signs or symptoms of infection and treat appropriately, including interruption or discontinuation of BLINCYTO[®] as needed.
- **Tumor Lysis Syndrome (TLS),** which may be life-threatening or fatal, has been observed. Preventive measures, including

pretreatment nontoxic cytoreduction and on-treatment hydration, should be used during BLINCYTO[®] treatment. Monitor patients for signs and symptoms of TLS and interrupt or discontinue BLINCYTO[®] as needed to manage these events.

- Neutropenia and Febrile Neutropenia, including life-threatening cases, have been observed. Monitor appropriate laboratory parameters (including, but not limited to, white blood cell count and absolute neutrophil count) during BLINCYTO[®] infusion and interrupt BLINCYTO[®] if prolonged neutropenia occurs.
- Effects on Ability to Drive and Use Machines: Due to the possibility of neurological events, including seizures, patients receiving BLINCYTO[®] are at risk for loss of consciousness, and should be advised against driving and engaging in hazardous occupations or activities such as operating heavy or potentially dangerous machinery while BLINCYTO[®] is being administered.
- Elevated Liver Enzymes: Transient elevations in liver enzymes have been associated with BLINCYTO[®] treatment with a median time to onset of 3 days. In patients receiving BLINCYTO[®], although the majority of these events were observed in the setting of CRS, some cases of elevated liver enzymes were observed outside the setting of CRS, with a median time to onset of 19 days. Grade 3 or greater elevations in liver enzymes occurred in approximately 7% of patients outside the setting of CRS and resulted in treatment discontinuation in less than 1% of patients. Monitor ALT, AST, gamma-glutamyl transferase (GGT), and TBIL prior to the start of and during BLINCYTO[®] treatment. BLINCYTO[®] treatment should be interrupted if transaminases rise to > 5 times the upper limit of normal (ULN) or if TBIL rises to > 3 times ULN.
- Pancreatitis: Fatal pancreatitis has been reported in patients receiving BLINCYTO[®] in combination with dexamethasone in clinical trials and the post-marketing setting. Evaluate patients who develop signs and symptoms of pancreatitis and interrupt or discontinue BLINCYTO[®] and dexamethasone as needed.
- Leukoencephalopathy: Although the clinical significance is unknown, cranial magnetic resonance imaging (MRI) changes showing leukoencephalopathy have been observed in patients receiving BLINCYTO[®], especially in patients previously treated with cranial irradiation and antileukemic chemotherapy.
- Preparation and administration errors have occurred with BLINCYTO[®] treatment. Follow instructions for preparation (including admixing) and administration in the PI strictly to minimize medication errors (including underdose and overdose).
- Immunization: Vaccination with live virus vaccines is not recommended for at least 2 weeks prior to the start of BLINCYTO[®] treatment, during treatment, and until immune recovery following last cycle of BLINCYTO[®].
- Risk of Serious Adverse Reactions in Pediatric Patients due to Benzyl Alcohol Preservative: Serious and fatal adverse reactions including "gasping syndrome," which is characterized by central nervous system depression, metabolic acidosis, and gasping respirations, can occur in neonates and infants treated with benzyl alcohol-preserved drugs including BLINCYTO[®] (with preservative). When prescribing BLINCYTO[®] (with preservative) for pediatric patients, consider the combined daily metabolic load of benzyl alcohol from all sources including BLINCYTO[®] (with preservative) and other drugs containing benzyl alcohol. The minimum amount of benzyl alcohol at which serious adverse reactions may occur is not known. Due to the addition of bacteriostatic saline, 7-day bags of BLINCYTO[®] solution for infusion with preservative contain benzyl alcohol and are not recommended for use in any patients weighing < 22 kg.

Adverse Reactions

- The most common adverse reactions ($\geq 20\%$) in clinical trial experience of patients with MRD-positive B-cell precursor ALL (BLAST Study) treated with BLINCYTO[®] were pyrexia, infusion related reactions, headache, infections (pathogen unspecified), tremor, and chills. Serious adverse reactions were reported in 61% of patients. The most common serious adverse reactions ($\geq 2\%$) included pyrexia, tremor, encephalopathy, aphasia, lymphopenia, neutropenia, overdose, device related infection, seizure, and staphylococcal infection.
- The most common adverse reactions ($\geq 20\%$) in clinical trial experience of patients with Philadelphia chromosome-negative relapsed or refractory B-cell precursor ALL (TOWER Study) treated with BLINCYTO[®] were infections (bacterial and pathogen unspecified), pyrexia, headache, infusion-related reactions, anemia, febrile neutropenia, thrombocytopenia, and neutropenia. Serious adverse reactions were reported in 62% of patients. The most common serious adverse reactions ($\geq 2\%$) included febrile neutropenia, pyrexia, sepsis, pneumonia, overdose, septic shock, CRS, bacterial sepsis, device related infection, and bacteremia.
- Adverse reactions that were observed more frequently ($\geq 10\%$) in the pediatric population compared to the adult population were pyrexia (80% vs. 61%), hypertension (26% vs. 8%), anemia (41% vs. 24%), infusion-related reaction (49% vs. 34%), thrombocytopenia (34% vs. 21%), leukopenia (24% vs. 11%), and weight increase (17% vs. 6%).
- In pediatric patients less than 2 years old (infants), the incidence of neurologic toxicities was not significantly different than for the other age groups, but its manifestations were different; the only event terms reported were agitation, headache, insomnia, somnolence, and irritability. Infants also had an increased incidence of hypokalemia (50%) compared to other pediatric age cohorts (15-20%) or adults (17%).

Dosage and Administration Guidelines

- BLINCYTO[®] is administered as a continuous intravenous infusion at a constant flow rate using an infusion pump which

should be programmable, lockable, non-elastomeric, and have an alarm.

- It is very important that the instructions for preparation (including admixing) and administration provided in the full Prescribing Information are strictly followed to minimize medication errors (including underdose and overdose).

Please see full Prescribing Information, including Boxed WARNINGS and Medication Guide, for BLINCYTO®.

About Amgen's Commitment to Oncology

Amgen Oncology is committed to helping patients take on some of the toughest cancers, such as those that have been resistant to drugs, those that progress rapidly through the body and those where limited treatment options exist. Amgen's supportive care treatments help patients combat certain side effects of strong chemotherapy, and our targeted medicines and immunotherapies focus on more than a dozen different malignancies, ranging from blood cancers to solid tumors. With decades of experience providing therapies for cancer patients, Amgen continues to grow its portfolio of innovative and biosimilar oncology medicines.

For more information, follow us on www.twitter.com/amgenoncology.

About Amgen

Amgen is committed to unlocking the potential of biology for patients suffering from serious illnesses by discovering, developing, manufacturing and delivering innovative human therapeutics. This approach begins by using tools like advanced human genetics to unravel the complexities of disease and understand the fundamentals of human biology.

Amgen focuses on areas of high unmet medical need and leverages its expertise to strive for solutions that improve health outcomes and dramatically improve people's lives. A biotechnology pioneer since 1980, Amgen has grown to be one of the world's leading independent biotechnology companies, has reached millions of patients around the world and is developing a pipeline of medicines with breakaway potential.

For more information, visit www.amgen.com and follow us on www.twitter.com/amgen.

Forward-Looking Statements

This news release contains forward-looking statements that are based on the current expectations and beliefs of Amgen. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including estimates of revenues, operating margins, capital expenditures, cash, other financial metrics, expected legal, arbitration, political, regulatory or clinical results or practices, customer and prescriber patterns or practices, reimbursement activities and outcomes and other such estimates and results. Forward-looking statements involve significant risks and uncertainties, including those discussed below and more fully described in the Securities and Exchange Commission reports filed by Amgen, including our most recent annual report on Form 10-K and any subsequent periodic reports on Form 10-Q and current reports on Form 8-K. Unless otherwise noted, Amgen is providing this information as of the date of this news release and does not undertake any obligation to update any forward-looking statements contained in this document as a result of new information, future events or otherwise.

No forward-looking statement can be guaranteed and actual results may differ materially from those we project. Discovery or identification of new product candidates or development of new indications for existing products cannot be guaranteed and movement from concept to product is uncertain; consequently, there can be no guarantee that any particular product candidate or development of a new indication for an existing product will be successful and become a commercial product. Further, preclinical results do not guarantee safe and effective performance of product candidates in humans. The complexity of the human body cannot be perfectly, or sometimes, even adequately modeled by computer or cell culture systems or animal models. The length of time that it takes for us to complete clinical trials and obtain regulatory approval for product marketing has in the past varied and we expect similar variability in the future. Even when clinical trials are successful, regulatory authorities may question the sufficiency for approval of the trial endpoints we have selected. We develop product candidates internally and through licensing collaborations, partnerships and joint ventures. Product candidates that are derived from relationships may be subject to disputes between the parties or may prove to be not as effective or as safe as we may have believed at the time of entering into such relationship. Also, we or others could identify safety, side effects or manufacturing problems with our products, including our devices, after they are on the market.

Our results may be affected by our ability to successfully market both new and existing products domestically and internationally, clinical and regulatory developments involving current and future products, sales growth of recently launched products, competition from other products including biosimilars, difficulties or delays in manufacturing our products and global economic conditions. In addition, sales of our products are affected by pricing pressure, political and public scrutiny and reimbursement policies imposed by third-party payers, including governments, private insurance plans and managed care providers and may be affected by regulatory, clinical and guideline developments and domestic and international trends toward managed care and healthcare cost containment. Furthermore, our research, testing, pricing, marketing and other operations are subject to extensive regulation by domestic and foreign government regulatory authorities. Our business may be impacted by government investigations, litigation and product liability claims. In addition, our business may be impacted by the adoption of new tax legislation or exposure to additional tax liabilities. While we routinely obtain patents for our products and technology, the protection offered by our patents and patent applications may be challenged, invalidated or circumvented by our competitors, or we may fail to prevail in present and future intellectual property litigation. We perform a substantial amount of our commercial manufacturing activities at a few key facilities, including in Puerto Rico, and also depend on third parties for a portion of our manufacturing activities, and limits on supply may constrain sales of certain of our current products and product candidate development. In addition, we compete with other companies with respect to many of our marketed products as well as for the discovery and development of new products. Further, some raw materials, medical devices and component parts for our products are supplied by sole third-party suppliers. Certain of our distributors, customers and payers have substantial purchasing leverage in their dealings with us. The discovery of significant problems with a product similar to one of our products that implicate an entire class of products could have a material adverse effect on sales of the affected products and on our business and results of operations. Our efforts to acquire other companies or products and to integrate the operations of companies we have acquired may not be successful. A breakdown, cyberattack or information security breach could compromise the confidentiality, integrity and availability of our systems and our data. Our stock price is volatile and may be affected by a number of events. Our business performance could affect or limit the ability of our Board of Directors to declare a dividend or our ability to pay a dividend or repurchase our common stock. We may not be able to access the capital and credit markets on terms that are favorable to us, or at all.

CONTACT: Amgen, Thousand Oaks
Kristen Davis, 805-447-3008 (Media)
Trish Hawkins, 805-447-5631 (Media)


Arvind Sood, 805-447-1060 (Investors)

Amgen, Europe
Emma Gilbert, +41 413692542

References:

1. Paeitta E. Assessing minimal residual disease (MRD) in leukemia: a changing definition and concept? *Bone Marrow Transplant.* 2002;29:459-465
2. Gökbuget N, et al. Adult patients with acute lymphoblastic leukemia and molecular failure display a poor prognosis and are candidates for stem cell transplantation and targeted therapies. *Blood.* 2012;120:1868-1876.
3. Brüggemann M, et al. Has MRD monitoring superseded other prognostic factors in adult ALL? *Blood.* 2012;120:4470-4481.
4. Cancer Research UK. About acute lymphoblastic leukaemia (ALL). <http://www.cancerresearchuk.org/about-cancer/acute-lymphoblastic-leukaemia-all/about>. Accessed Nov. 1, 2018.
5. Mayo Clinic. Acute lymphocytic leukemia. <http://www.mayoclinic.org/diseases-conditions/acute-lymphocytic-leukemia/basics/definition/con-20042915>. Accessed Nov. 1, 2018.
6. Bassan R, Spinelli O, Oldani E, et al. Improved risk classification for risk-specific therapy based on the molecular study of minimal residual disease (MRD) in adult acute lymphoblastic leukemia (ALL). *Blood.* 2009;113: 4153-4162.

The logo for Amgen, featuring the word "AMGEN" in a bold, blue, sans-serif font. A registered trademark symbol (®) is located at the top right of the letter "N".

 View original content to download multimedia: <http://www.prnewswire.com/news-releases/european-commission-approves-blincyto-blinatumomab-in-patients-with-philadelphia-chromosome-negative-minimal-residual-disease-positive-b-cell-precursor-acute-lymphoblastic-leukemia-300781842.html>

SOURCE Amgen