



Amgen Announces Positive Top-Line Results From Otezla® (apremilast) Phase 3 ADVANCE Study In Mild-To-Moderate Plaque Psoriasis

May 6, 2020

Otezla Treatment Resulted in Significant Improvements in Measures of Mild-to-Moderate Psoriasis Compared With Placebo

Data Will be Submitted to the U.S. Food and Drug Administration (FDA) for Inclusion in Otezla Prescribing Information

THOUSAND OAKS, Calif., May 6, 2020 /PRNewswire/ -- Amgen (NASDAQ:AMGN) today announced positive top-line results from the ADVANCE trial, a Phase 3, multicenter, randomized, placebo-controlled, double-blind study to assess the efficacy of Otezla® (apremilast) in adults with mild-to-moderate plaque psoriasis. The study showed that oral Otezla 30 mg twice daily achieved a statistically significant improvement, compared with placebo, in the primary endpoint of the static Physician's Global Assessment (sPGA) response (defined as an sPGA score of clear (0) or almost clear (1) with at least a 2-point reduction from baseline) at week 16.

In addition, the week 16 secondary endpoints of achieving at least 75% improvement from baseline in the percent of affected body surface area (BSA); change in BSA total score from baseline; and change in Psoriasis Area and Severity Index (PASI) total score from baseline were each also statistically significant for the treatment effect of Otezla compared with placebo.

"Many patients with mild-to-moderate plaque psoriasis who use topical therapies still have challenges managing their psoriasis," said David M. Reese, M.D., executive vice president of Research and Development at Amgen. "We look forward to discussions with the FDA about the potential to bring Otezla, which has already been prescribed to hundreds of thousands of patients with moderate-to-severe psoriasis, to more patients who may need additional therapeutic options."

The adverse events observed in this trial were consistent with the known safety profile of Otezla. The most commonly reported adverse events that occurred in at least 5% of patients in either treatment group were diarrhea, headache, nausea, nasopharyngitis and upper respiratory tract infection.

Detailed results will be submitted for presentation at an upcoming medical meeting.

In the U.S., Otezla is approved for the treatment of adult patients with moderate-to-severe plaque psoriasis who are candidates for phototherapy or systemic therapy, adult patients with active psoriatic arthritis and for adult patients with oral ulcers associated with Behçet's Disease. Since its initial FDA approval in 2014, Otezla has been prescribed to more than 250,000 patients with moderate-to-severe plaque psoriasis or active psoriatic arthritis in the U.S.^[1]

About ADVANCE (PSOR-022)

ADVANCE (PSOR-022) is a Phase 3, multicenter, randomized, placebo-controlled, double-blind study evaluating the efficacy and safety of Otezla in patients with mild-to-moderate plaque psoriasis (defined as BSA involvement of 2% to 15%, PASI score of 2 to 15, sPGA score of 2 to 3). The study randomized 595 patients 1:1 to receive Otezla 30 mg twice daily or placebo for the first 16 weeks. All patients then received Otezla during an open-label extension phase through week 32.

The primary endpoint was the percentage of patients with sPGA response [defined as a sPGA score of clear (0) or almost clear (1) with at least a 2-point reduction from baseline] at week 16.

About Psoriasis

Psoriasis is a serious, chronic inflammatory disease that causes raised, red, scaly patches to appear on the skin, typically affecting the outside of the elbows, knees or scalp, though it can appear on any location.^[2] Approximately 125 million people worldwide have psoriasis, including around 14 million people in Europe and more than 7.5 million people in the United States.^{[3],[4]} About 80% of those patients have plaque psoriasis.^[5]

About Otezla® (apremilast)

Otezla® (apremilast) 30 mg tablets is an oral small-molecule inhibitor of phosphodiesterase 4 (PDE4) specific for cyclic adenosine monophosphate (cAMP). PDE4 inhibition results in increased intracellular cAMP levels, which is thought to indirectly modulate the production of inflammatory mediators. The specific mechanism(s) by which Otezla exerts its therapeutic action in patients is not well defined.

Otezla® (apremilast) U.S. INDICATIONS

Otezla® (apremilast) is indicated for the treatment of adult patients with moderate to severe plaque psoriasis who are candidates for phototherapy or systemic therapy.

Otezla is indicated for the treatment of adult patients with active psoriatic arthritis.

Otezla is indicated for the treatment of adult patients with oral ulcers associated with Behçet's Disease.

Otezla® (apremilast) U.S. IMPORTANT SAFETY INFORMATION

Contraindications

- Otezla® (apremilast) is contraindicated in patients with a known hypersensitivity to apremilast or to any of the excipients in

the formulation

Warnings and Precautions

- **Diarrhea, Nausea, and Vomiting:** Cases of severe diarrhea, nausea, and vomiting were associated with the use of Otezla. Most events occurred within the first few weeks of treatment. In some cases patients were hospitalized. Patients 65 years of age or older and patients taking medications that can lead to volume depletion or hypotension may be at a higher risk of complications from severe diarrhea, nausea, or vomiting. Monitor patients who are more susceptible to complications of diarrhea or vomiting; advise patients to contact their healthcare provider. Consider Otezla dose reduction or suspension if patients develop severe diarrhea, nausea, or vomiting
- **Depression:** Carefully weigh the risks and benefits of treatment with Otezla for patients with a history of depression and/or suicidal thoughts/behavior, or in patients who develop such symptoms while on Otezla. Patients, caregivers, and families should be advised of the need to be alert for the emergence or worsening of depression, suicidal thoughts or other mood changes, and they should contact their healthcare provider if such changes occur
 - **Psoriasis:** Treatment with Otezla is associated with an increase in depression. During clinical trials, 1.3% (12/920) of patients reported depression compared to 0.4% (2/506) on placebo. Depression was reported as serious in 0.1% (1/1308) of patients exposed to Otezla, compared to none in placebo-treated patients (0/506). Suicidal behavior was observed in 0.1% (1/1308) of patients on Otezla, compared to 0.2% (1/506) on placebo. One patient treated with Otezla attempted suicide; one patient on placebo committed suicide
 - **Psoriatic Arthritis:** Treatment with Otezla is associated with an increase in depression. During clinical trials, 1.0% (10/998) reported depression or depressed mood compared to 0.8% (4/495) treated with placebo. Suicidal ideation and behavior was observed in 0.2% (3/1441) of patients on Otezla, compared to none in placebo-treated patients. Depression was reported as serious in 0.2% (3/1441) of patients exposed to Otezla, compared to none in placebo-treated patients (0/495). Two patients who received placebo committed suicide compared to none on Otezla
 - **Behçet's Disease:** Treatment with Otezla is associated with an increase in depression. During the clinical trial, 1% (1/104) reported depression or depressed mood compared to 1% (1/103) treated with placebo. No instances of suicidal ideation or behavior were reported in patients treated with Otezla or treated with placebo
- **Weight Decrease:** Monitor body weight regularly; evaluate unexplained or clinically significant weight loss, and consider discontinuation of Otezla
 - **Psoriasis:** Body weight loss of 5-10% occurred in 12% (96/784) of patients treated with Otezla and in 5% (19/382) of patients treated with placebo. Body weight loss of \geq 10% occurred in 2% (16/784) of patients treated with Otezla compared to 1% (3/382) of patients treated with placebo
 - **Psoriatic Arthritis:** Body weight loss of 5-10% was reported in 10% (49/497) of patients taking Otezla and in 3.3% (16/495) of patients taking placebo
 - **Behçet's Disease:** Body weight loss of $>$ 5% was reported in 4.9% (5/103) of patients taking Otezla and in 3.9% (4/102) of patients taking placebo
- **Drug Interactions:** Apremilast exposure was decreased when Otezla was co-administered with rifampin, a strong CYP450 enzyme inducer; loss of Otezla efficacy may occur. Concomitant use of Otezla with CYP450 enzyme inducers (e.g., rifampin, phenobarbital, carbamazepine, phenytoin) is not recommended

Adverse Reactions

- **Psoriasis:** Adverse reactions reported in \geq 5% of patients were (Otezla%, placebo%): diarrhea (17, 6), nausea (17, 7), upper respiratory tract infection (9, 6), tension headache (8, 4), and headache (6, 4)
- **Psoriatic Arthritis:** Adverse reactions reported in at least 2% of patients taking Otezla, that occurred at a frequency at least 1% higher than that observed in patients taking placebo, for up to 16 weeks (after the initial 5-day titration), were (Otezla%, placebo%): diarrhea (7.7, 1.6); nausea (8.9, 3.1); headache (5.9, 2.2); upper respiratory tract infection (3.9, 1.8); vomiting (3.2, 0.4); nasopharyngitis (2.6, 1.6); upper abdominal pain (2.0, 0.2)
- **Behçet's Disease:** Adverse reactions reported in at least \geq 5% of patients taking Otezla, that occurred at a frequency at least 1% higher than that observed in patients taking placebo, for up to 12 weeks, were (Otezla%, placebo%): diarrhea (41.3, 20.4); nausea (19.2, 10.7); headache (14.4, 10.7); upper respiratory tract infection (11.5, 4.9); upper abdominal pain (8.7, 1.9); vomiting (8.7, 1.9); back pain (7.7, 5.8); viral upper respiratory tract infection (6.7, 4.9); arthralgia (5.8, 2.9)

Use in Specific Populations

- **Pregnancy:** Otezla has not been studied in pregnant women. Advise pregnant women of the potential risk of fetal loss. Consider pregnancy planning and prevention for females of reproductive potential. There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to Otezla during pregnancy. Information about the registry can be obtained by calling 1-877-311-8972 or visiting <https://mothertobaby.org/ongoing-study/otezla/>
- **Lactation:** There are no data on the presence of apremilast or its metabolites in human milk, the effects of apremilast on the breastfed infant, or the effects of the drug on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Otezla and any potential adverse effects on the breastfed

child from Otezla or from the underlying maternal condition

- Renal Impairment: Otezla dosage should be reduced in patients with severe renal impairment (creatinine clearance less than 30 mL/min) for details, see Dosage and Administration, Section 2, in the Full Prescribing Information

Please click [here](#) for Otezla® Full Prescribing Information.

About Amgen

Amgen is committed to unlocking the potential of biology for patients suffering from serious illnesses by discovering, developing, manufacturing and delivering innovative human therapeutics. This approach begins by using tools like advanced human genetics to unravel the complexities of disease and understand the fundamentals of human biology.

Amgen focuses on areas of high unmet medical need and leverages its expertise to strive for solutions that improve health outcomes and dramatically improve people's lives. A biotechnology pioneer since 1980, Amgen has grown to be one of the world's leading independent biotechnology companies, has reached millions of patients around the world and is developing a pipeline of medicines with breakaway potential.

For more information, visit www.amgen.com and follow us on www.twitter.com/amgen.

Forward-Looking Statements

This news release contains forward-looking statements that are based on the current expectations and beliefs of Amgen. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including any statements on the outcome, benefits and synergies of collaborations, or potential collaborations, with any other company, including Adaptive Biotechnologies (including statements regarding such collaboration's ability to discover and develop fully-human neutralizing antibodies targeting SARS-CoV-2 to potentially prevent or treat COVID-19), BeiGene, Ltd., or the Otezla® (apremilast) acquisition, including anticipated Otezla sales growth and the timing of non-GAAP EPS accretion, as well as estimates of revenues, operating margins, capital expenditures, cash, other financial metrics, expected legal, arbitration, political, regulatory or clinical results or practices, customer and prescriber patterns or practices, reimbursement activities and outcomes, effects of pandemics or other widespread health problems such as the ongoing COVID-19 pandemic on our business, outcomes, progress, or effects relating to studies of Otezla as a potential treatment for COVID-19, and other such estimates and results. Forward-looking statements involve significant risks and uncertainties, including those discussed below and more fully described in the Securities and Exchange Commission reports filed by Amgen, including our most recent annual report on Form 10-K and any subsequent periodic reports on Form 10-Q and current reports on Form 8-K. Unless otherwise noted, Amgen is providing this information as of the date of this news release and does not undertake any obligation to update any forward-looking statements contained in this document as a result of new information, future events or otherwise.

No forward-looking statement can be guaranteed and actual results may differ materially from those we project. Discovery or identification of new product candidates or development of new indications for existing products cannot be guaranteed and movement from concept to product is uncertain; consequently, there can be no guarantee that any particular product candidate or development of a new indication for an existing product will be successful and become a commercial product. Further, preclinical results do not guarantee safe and effective performance of product candidates in humans. The complexity of the human body cannot be perfectly, or sometimes, even adequately modeled by computer or cell culture systems or animal models. The length of time that it takes for us to complete clinical trials and obtain regulatory approval for product marketing has in the past varied and we expect similar variability in the future. Even when clinical trials are successful, regulatory authorities may question the sufficiency for approval of the trial endpoints we have selected. We develop product candidates internally and through licensing collaborations, partnerships and joint ventures. Product candidates that are derived from relationships may be subject to disputes between the parties or may prove to be not as effective or as safe as we may have believed at the time of entering into such relationship. Also, we or others could identify safety, side effects or manufacturing problems with our products, including our devices, after they are on the market.

Our results may be affected by our ability to successfully market both new and existing products domestically and internationally, clinical and regulatory developments involving current and future products, sales growth of recently launched products, competition from other products including biosimilars, difficulties or delays in manufacturing our products and global economic conditions. In addition, sales of our products are affected by pricing pressure, political and public scrutiny and reimbursement policies imposed by third-party payers, including governments, private insurance plans and managed care providers and may be affected by regulatory, clinical and guideline developments and domestic and international trends toward managed care and healthcare cost containment. Furthermore, our research, testing, pricing, marketing and other operations are subject to extensive regulation by domestic and foreign government regulatory authorities. Our business may be impacted by government investigations, litigation and product liability claims. In addition, our business may be impacted by the adoption of new tax legislation or exposure to additional tax liabilities. If we fail to meet the compliance obligations in the corporate integrity agreement between us and the U.S. government, we could become subject to significant sanctions. Further, while we routinely obtain patents for our products and technology, the protection offered by our patents and patent applications may be challenged, invalidated or circumvented by our competitors, or we may fail to prevail in present and future intellectual property litigation. We perform a substantial amount of our commercial manufacturing activities at a few key facilities, including in Puerto Rico, and also depend on third parties for a portion of our manufacturing activities, and limits on supply may constrain sales of certain of our current products and product candidate development. An outbreak of disease or similar public health threat, such as COVID-19, and the public and governmental effort to mitigate against the spread of such disease, could have a significant adverse effect on the supply of materials for our manufacturing activities, the distribution of our products, the commercialization of our product candidates, and our clinical trial operations, and any such events may have a material adverse effect on our product development, product sales, business and results of operations. We rely on collaborations with third parties for the development of some of our product candidates and for the commercialization and sales of some of our commercial products. In addition, we compete with other companies with respect to many of our marketed products as well as for the discovery and development of new products. Further, some raw materials, medical devices and component parts for our products are supplied by sole third-party suppliers. Certain of our distributors, customers and payers have substantial purchasing leverage in their dealings with us. The discovery of significant problems with a product similar to one of our products that implicate an entire class of products could have a material adverse effect on sales of the affected products and on our business and results of operations. Our efforts to collaborate with or acquire other companies, products or technology, and to integrate the operations of companies or to support the products or technology we have acquired, may not be successful. A breakdown, cyberattack or information security breach could compromise the confidentiality, integrity and availability of our systems and our data. Our stock price is volatile and may be affected by a number of events. Our business performance could affect or limit the ability of our Board of Directors to declare a dividend or our ability to pay a dividend or repurchase our common stock. We may not be able to access the capital and credit markets on terms that are favorable to us, or at all.

Any scientific information discussed in this news release relating to new indications for our products is preliminary and investigative and is not part of the labeling approved by the U.S. Food and Drug Administration for the products. The products are not approved for the investigational use(s) discussed in this news release, and no conclusions can or should be drawn regarding the safety or effectiveness of the products for these uses.

CONTACT: Amgen, Thousand Oaks
Megan Fox, 805-447-1423 (media)
Trish Rowland, 805-447-5631 (media)
Arvind Sood, 805-447-1060 (investors)



[1] Data on File at Amgen Inc.

[2] National Psoriasis Foundation. About Psoriasis. Available at: <https://www.psoriasis.org/about-psoriasis>. Accessed January 24, 2020.

[3] National Psoriasis Foundation. Statistics. Available at: <https://www.psoriasis.org/content/statistics>. Accessed January 24, 2020.

[4] Ortonne JP, Prinz JC. Alefacept: a novel and selective biologic agent for the treatment of chronic plaque psoriasis. *Eur J Dermatol*. 2004;14(1):41–45.

[5] National Psoriasis Foundation. Plaque Psoriasis. Available at: <https://www.psoriasis.org/about-psoriasis/types/plaque>. Accessed January 24, 2020.

 View original content to download multimedia: <http://www.prnewswire.com/news-releases/amgen-announces-positive-top-line-results-from-otezla-apremilast-phase-3-advance-study-in-mild-to-moderate-plaque-psoriasis-301053621.html>

SOURCE Amgen