



IMDELLTRA® IN COMBINATION WITH IMFINZI® DEMONSTRATED LANDMARK IMPROVEMENT IN OVERALL SURVIVAL IN FIRST-LINE EXTENSIVE STAGE SMALL CELL LUNG CANCER

September 8, 2026

Phase 3 DeLLphi-305 Study Delivered Highly Significant and Clinically Meaningful Overall Survival Benefit Versus Durvalumab Alone Study Also Met Secondary Progression-Free Survival and Objective Response Rate Endpoints

THOUSAND OAKS, Calif., Sept. 8, 2026 /PRNewswire/ -- Amgen (NASDAQ: AMGN) today announced landmark results from the Phase 3 DeLLphi-305 study, which met its primary endpoint at a pre-specified interim analysis, demonstrating a statistically significant and clinically meaningful improvement in overall survival with IMDELLTRA® (tarlatamab-dlle) in combination with AstraZeneca's Imfinzi® (durvalumab) compared to durvalumab alone. This study tested IMDELLTRA as a first-line maintenance treatment for people with extensive stage small cell lung cancer (ES-SCLC) whose disease had not progressed following initial treatment with durvalumab, platinum-based chemotherapy and etoposide. The study also demonstrated a statistically significant and clinically meaningful improvement in progression-free survival (PFS) and objective response rate (ORR) among patients treated with IMDELLTRA in combination with durvalumab.

Overall, the safety profile of IMDELLTRA in combination with durvalumab was consistent with the known safety profiles of the individual medicines. No new or unexpected safety signals were identified in DeLLphi-305. The study design included monitoring in a healthcare setting for 1 to 2 hours (6 to 8 hours in certain regions including Europe) following IMDELLTRA infusion on Cycle 1 Day 1 and Cycle 1 Day 8.

"IMDELLTRA has already revolutionized the standard of survival for patients with extensive stage small cell lung cancer whose disease progressed on or after prior treatment. These landmark results from DeLLphi-305 suggest IMDELLTRA will further revolutionize the standard for survival earlier in the treatment journey and meaningfully shift the treatment paradigm for people facing this devastating disease," said Jay Bradner, M.D., executive vice president, Research and Development, Artificial Intelligence and Data at Amgen. "Despite clinical advances, extensive stage small cell lung cancer remains one of the most aggressive and difficult-to-treat cancers, and substantial improvements in survival are rare, underscoring the importance of these findings."

This marks the first Phase 3 study including a Bispecific T-cell Engager (BiTE®) therapy to demonstrate an overall survival benefit in the first-line maintenance setting for ES-SCLC.

"Given the aggressive nature of small cell lung cancer, many patients quickly relapse on current therapy and never reach second-line treatment. These patients do not have time to wait, making substantial progress in the first-line setting critically important," said Jacob Sands, MD, Associate Chief of the Lowe Center for Thoracic Oncology at Dana-Farber Cancer Institute. "In my career treating people with extensive stage small cell lung cancer, these are among the most compelling survival results I have seen, indicating the potential to reshape the natural history of small cell lung cancer. DeLLphi-305 represents an unprecedented milestone and suggests we may be entering a new era where meaningfully longer survival is possible for more patients."

Following the U.S. Food and Drug Administration (FDA) approval of IMDELLTRA for adults with ES-SCLC whose disease has progressed on or after platinum-based chemotherapy, the robust global DeLLphi clinical trial program has continued to evaluate its potential across earlier stages of disease and lines of treatment, both as a monotherapy and in combination regimens. Building on the DeLLphi-303 trial, which showed very encouraging efficacy and tolerability results for IMDELLTRA in combination with anti-PD-L1 checkpoint inhibitors in the treatment of first-line ES-SCLC, DeLLphi-305 was designed to evaluate IMDELLTRA earlier in the course of disease, where more patients with SCLC may have the opportunity to benefit.

SCLC is one of the most aggressive forms of lung cancer, accounting for approximately 13-15% of the more than 2.6 million lung cancer cases diagnosed worldwide each year.¹⁻³ Despite advances in the treatment of ES-SCLC, outcomes remain poor, with a median survival of approximately one year from the start of first-line maintenance treatment and only about 40% of patients receiving second-line therapy.⁴⁻⁷

Amgen plans to present detailed data from DeLLphi-305 at an upcoming international medical congress and share it with regulatory authorities.

About the Phase 3 DeLLphi-305 Study

The DeLLphi-305 trial is sponsored by Amgen, with partial funding and durvalumab provided by AstraZeneca. It is a global Phase 3, randomized, open-label clinical trial evaluating the efficacy and safety of IMDELLTRA in combination with durvalumab compared to durvalumab alone as first-line maintenance treatment for patients with ES-SCLC who had not progressed following treatment with durvalumab, platinum-based chemotherapy and etoposide. Five hundred and sixty-three patients who completed initial treatment with durvalumab, platinum-based chemotherapy and etoposide were randomized 1:1 to receive either IMDELLTRA in combination with durvalumab or durvalumab alone until progression or unacceptable toxicity. Following IMDELLTRA infusion on Cycle 1 Day 1 and Cycle 1 Day 8, patients were monitored in a healthcare setting for 1 to 2 hours (6 to 8 hours in certain regions including Europe). The trial included patients with both treated and untreated asymptomatic brain metastases at baseline. The primary outcome measure of the trial is OS. Secondary endpoints include progression-free survival (PFS) and objective response rate (ORR).⁸

About Tarlatamab Clinical Trials

Tarlatamab is being investigated in multiple studies including DeLLphi-303, a Phase 1b study investigating tarlatamab in combination with standard-of-care (SOC) therapies in first-line ES-SCLC; DeLLphi-304, a randomized Phase 3 study comparing tarlatamab monotherapy with SOC chemotherapy in second-line treatment of SCLC; DeLLphi-305, a randomized Phase 3 study comparing tarlatamab in combination with durvalumab vs. durvalumab alone as first-line maintenance treatment in ES-SCLC; DeLLphi-306, a randomized placebo-controlled Phase 3 study of tarlatamab following concurrent chemoradiotherapy in limited-stage SCLC; DeLLphi-308, a Phase 1b study evaluating subcutaneous tarlatamab in second-line or later ES-SCLC; DeLLphi-309, a Phase 2 study evaluating alternative intravenous dosing regimens with tarlatamab in second-line ES-SCLC; DeLLphi-310, a Phase 1b study of tarlatamab in combination with YL201 with or without durvalumab in patients with ES-SCLC; DeLLphi-311, a Phase

1b study of tarlatamab in combination with etakafusp alfa (AB248), a novel CD8+ T-cell selective interleukin-2 (IL-2), in patients with ES-SCLC; DeLLphi-312, a randomized Phase 3 study evaluating tarlatamab in combination with durvalumab, carboplatin and etoposide as an induction and maintenance therapy in first-line treatment of ES-SCLC; DeLLphi-313, a Phase 1b study of tarlatamab in combination with zocilurtatug pelitecan, a DLL3 targeting antibody drug conjugate, with and without durvalumab in patients with ES-SCLC, and DeLLphi-315, a Phase 3 study of subcutaneous tarlatamab in patients with second-line ES-SCLC.⁹

For more information, please visit www.tarlatamabclinicaltrials.com.

About IMDELLTRA® (tarlatamab-dlle)

IMDELLTRA is a first-in-class targeted immunotherapy engineered by Amgen researchers to bind to both DLL3 on tumor cells and CD3 on T cells, thereby activating T cells to kill DLL3-expressing SCLC cells. This results in the formation of a cytolytic synapse with lysis of the cancer cell.^{10,11} DLL3 is a protein that is expressed on the surface of SCLC cells in ~85-96% of patients with SCLC, but is minimally expressed on healthy cells, making it an exciting target.^{12,13}

U.S. INDICATION

IMDELLTRA® (tarlatamab-dlle) is indicated for the treatment of adult patients with extensive stage small cell lung cancer (ES-SCLC) with disease progression on or after platinum-based chemotherapy.

IMDELLTRA IMPORTANT SAFETY INFORMATION

WARNING: CYTOKINE RELEASE SYNDROME and NEUROLOGIC TOXICITY including IMMUNE EFFECTOR CELL-ASSOCIATED NEUROTOXICITY SYNDROME

- **Cytokine release syndrome (CRS), including life-threatening or fatal reactions, can occur in patients receiving IMDELLTRA®. Initiate treatment with IMDELLTRA® using the step-up dosing schedule to reduce the incidence and severity of CRS. Withhold IMDELLTRA® until CRS resolves or permanently discontinue based on severity.**
- **Neurologic toxicity and immune effector cell-associated neurotoxicity syndrome (ICANS), including life-threatening or fatal reactions, can occur in patients receiving IMDELLTRA®. Monitor patients for signs and symptoms of neurologic toxicity, including ICANS, during treatment and treat promptly. Withhold IMDELLTRA® until ICANS resolves or permanently discontinue based on severity.**

WARNINGS AND PRECAUTIONS

- **Cytokine Release Syndrome (CRS):** IMDELLTRA® can cause CRS including life-threatening or fatal reactions. In the pooled safety population, CRS occurred in 57% (268/473) of patients who received IMDELLTRA®, including 39% Grade 1, 15% Grade 2, 1.7% Grade 3 and 0.2% Grade 4. Recurrent CRS occurred in 24% of IMDELLTRA®-treated patients including 20% Grade 1 and 3.4% Grade 2; one patient experienced recurrent Grade 3.

Among the 268 patients who experienced CRS, 73% had CRS after the first dose, 60% had CRS after the second dose, and 15% had CRS following the third or later dose. Following the Cycle 1 Day 1, Day 8, Day 15 infusions, 24%, 8%, and 1% of patients experienced Grade ≥ 2 CRS, respectively. From Cycle 2 onwards, 1.5% of patients experienced Grade ≥ 2 CRS. Of the patients who experienced CRS, 31% received steroids and 10% required tocilizumab. The median time to onset of all grade CRS from most recent dose of IMDELLTRA® was 16 hours (range: start of infusion to 15 days). The median time to onset of Grade ≥ 2 CRS from most recent dose of IMDELLTRA® was 15 hours (range: start of infusion to 15 days).

Clinical signs and symptoms of CRS included pyrexia, hypotension, fatigue, tachycardia, headache, hypoxia, nausea, and vomiting. Potentially life-threatening complications of CRS may include cardiac dysfunction, acute respiratory distress syndrome, neurologic toxicity, renal and/or hepatic failure, and disseminated intravascular coagulation (DIC).

Administer IMDELLTRA® following the recommended step-up dosing and administer concomitant medications before and after Cycle 1 Day 1 and Cycle 1 Day 8 IMDELLTRA® infusions as described in Table 3 of the Prescribing Information (PI) to reduce the risk of CRS. Administer IMDELLTRA® in an appropriate healthcare facility equipped to monitor and manage CRS. Ensure patients are well hydrated prior to administration of IMDELLTRA®.

Closely monitor patients for signs and symptoms of CRS during treatment with IMDELLTRA®. At the first sign of CRS, immediately discontinue IMDELLTRA® infusion, evaluate the patient for hospitalization and institute supportive care based on severity. Withhold or permanently discontinue IMDELLTRA® based on severity. Counsel patients and caregivers to seek medical attention should signs or symptoms of CRS occur.

- **Neurologic Toxicity, Including ICANS:** IMDELLTRA® can cause life-threatening or fatal neurologic toxicity, including ICANS. In the pooled safety population, neurologic toxicity occurred in 65% of patients who received IMDELLTRA®, with

Grade 3 or higher events in 7% of patients including fatal events in 0.2%. The most frequent neurologic toxicities were dysgeusia (34%), headache (17%), peripheral neuropathy (9%), dizziness (9%), and insomnia (8%). The incidence of signs and symptoms consistent with ICANS was 10% in IMDELLTRA[®]-treated patients including events with the preferred terms: ICANS (4.7%), muscular weakness (3.2%), cognitive disorder (0.6%), aphasia (0.6%), depressed level of consciousness (0.4%), seizures (0.4%), encephalopathy (0.4%), and leukoencephalopathy (0.2%). There was one fatal reaction of ICANS. Recurrent ICANS occurred in 1.5% of patients. Of the patients who experienced ICANS, most experienced the event following Cycle 1 Day 1 (2.5%) and Cycle 1 Day 8 (3.6%). Following Day 1, Day 8, and Day 15 infusions, 1.3%, 1.3% and 0.4% of patients experienced Grade ≥ 2 ICANS, respectively. ICANS can occur several weeks following administration of IMDELLTRA[®]. The median time to onset of ICANS from the first dose of IMDELLTRA[®] was 16 days (range: 1 to 862 days). The median time to resolution of ICANS was 4 days (range: 1 to 40 days).

The onset of ICANS can be concurrent with CRS, following resolution of CRS, or in the absence of CRS. Clinical signs and symptoms of ICANS may include but are not limited to confusional state, depressed level of consciousness, disorientation, somnolence, lethargy, and bradyphrenia.

Patients receiving IMDELLTRA[®] are at risk of neurologic adverse reactions and ICANS resulting in depressed level of consciousness. Advise patients to refrain from driving and engaging in hazardous occupations or activities, such as operating heavy or potentially dangerous machinery, until neurologic symptoms resolve.

Closely monitor patients for signs and symptoms of neurologic toxicity and ICANS during treatment with IMDELLTRA[®]. At the first sign of ICANS, immediately discontinue the infusion, evaluate the patient and provide supportive therapy based on severity. Withhold IMDELLTRA[®] or permanently discontinue based on severity.

Cytopenias: IMDELLTRA[®] can cause cytopenias including neutropenia, thrombocytopenia, and anemia. In the pooled safety population, based on laboratory data, decreased neutrophils occurred in 16% of patients, including 9% Grade 3 or 4. The median time to onset for Grade 3 or 4 decreased neutrophil count was 41 days (range: 2 to 306 days). Decreased platelets occurred in 30% including 2.2% Grade 3 or 4. The median time to onset for Grade 3 or 4 decreased platelets was 67 days (range: 3 to 420 days). Decreased hemoglobin occurred in 56% of patients, including 4.7% Grade 3 or 4. Febrile neutropenia was reported as an adverse event in 1.5% of patients treated with IMDELLTRA[®].

Monitor patients for signs and symptoms of cytopenias. Perform complete blood counts prior to treatment with all doses of IMDELLTRA[®], up through Cycle 5 Day 15 and then prior to administration on Day 1 of each cycle starting with Cycle 6. Based on the severity of cytopenias, temporarily withhold, or permanently discontinue IMDELLTRA[®].

- **Infections:** IMDELLTRA[®] can cause serious infections, including life-threatening and fatal infections.

In the pooled safety population, infections, including opportunistic infections, occurred in 43% of patients who received IMDELLTRA[®], including 14% Grade 3 or 4. The most frequent infections were pneumonia (11%), urinary tract infection (9%), COVID-19 (6%), upper respiratory tract infection (4.7%), respiratory tract infection (4%), candida infection (2.1%), oral candidiasis (2.1%), and nasopharyngitis (2.1%).

Monitor patients for signs and symptoms of infection prior to and during treatment with IMDELLTRA[®] and treat as clinically indicated. Withhold or permanently discontinue IMDELLTRA[®] based on severity.

- **Hepatotoxicity:** IMDELLTRA[®] can cause hepatotoxicity. In the pooled safety population, based on laboratory data, elevated ALT occurred in 39% of patients who received IMDELLTRA[®], including 2.5% with Grade 3 or 4 ALT. Elevated AST occurred in 43% of patients, including 3.2% Grade 3 or 4. Elevated bilirubin also occurred in 16% of patients, including 1.3% Grade 3 or 4. Liver enzyme elevation can occur with or without concurrent CRS.

Monitor liver enzymes and bilirubin prior to treatment with IMDELLTRA[®], and as clinically indicated. Withhold IMDELLTRA[®] or permanently discontinue based on severity.

- **Hypersensitivity:** IMDELLTRA[®] can cause severe hypersensitivity reactions. Clinical signs and symptoms of hypersensitivity may include, but are not limited to, rash and bronchospasm. Monitor patients for signs and symptoms of hypersensitivity during treatment with IMDELLTRA[®] and manage as clinically indicated. Withhold or consider permanent discontinuation of IMDELLTRA[®] based on severity.
- **Embryo-Fetal Toxicity:** Based on its mechanism of action, IMDELLTRA[®] may cause fetal harm when administered to a

pregnant woman. Advise patients of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with IMDELLTRA® and for 2 months after the last dose.

ADVERSE REACTIONS

- The pooled safety population reflects exposure to intravenous IMDELLTRA®, as a single agent, at the recommended dosage of IMDELLTRA® 1 mg on Cycle 1 Day 1 followed by 10 mg on Days 8 and 15, and then every 2 weeks until disease progression or intolerable toxicity in 473 patients with small cell lung cancer enrolled in three clinical trials: DeLLphi-300, DeLLphi-301 and DeLLphi-304. Among 473 patients who received IMDELLTRA®, 40% were exposed for 6 months or longer and 19% were exposed for greater than one year.
- The most common (≥ 20%) adverse reactions were CRS (57%), fatigue (48%), decreased appetite (38%), dysgeusia (34%), pyrexia (33%), constipation (31%), musculoskeletal pain (31%) and nausea (25%).
- The most common (≥ 5%) Grade 3 or 4 laboratory abnormalities were decreased lymphocytes (43%), decreased sodium (12%), decreased total neutrophils (9%) and increased uric acid (6%).

DOSAGE AND ADMINISTRATION: Important Dosing Information

- Administer IMDELLTRA® as an intravenous infusion over 1 hour.
- Administer IMDELLTRA® according to the step-up dose and schedule in the IMDELLTRA® PI (Table 1) to reduce the incidence and severity of CRS.
- Evaluate complete blood count, liver enzymes and bilirubin prior to administration of all doses of IMDELLTRA® up through Cycle 5 Day 15 and then prior to administration of IMDELLTRA® on Day 1 of each cycle starting with Cycle 6. More frequent evaluation may be necessary if clinically indicated.
- For Cycle 1, administer recommended concomitant medications before and after Cycle 1 Day 1 and Cycle 1 Day 8 IMDELLTRA® infusions to reduce the risk of CRS reactions as described in the PI (Table 3).
- IMDELLTRA® should only be administered by a qualified healthcare professional with appropriate medical support to manage severe reactions such as CRS and neurologic toxicity including ICANS.
- Due to the risk of CRS and neurologic toxicity, including ICANS, monitor patients from the start of the IMDELLTRA® infusion for 22 to 24 hours following Cycle 1 Day 1 and Cycle 1 Day 8 in an appropriate healthcare setting.
- Recommend that patients remain within 1 hour of an appropriate healthcare setting for a total of 48 hours from the start of the infusion with IMDELLTRA® following Cycle 1 Day 1 and Cycle 1 Day 8 doses, accompanied by a caregiver.
- Inform both the patient and the caregiver on the signs and symptoms of CRS and ICANS prior to discharge.
- Ensure patients are well hydrated prior to administration of IMDELLTRA®.

Please see IMDELLTRA® [full Prescribing Information](#), including **BOXED WARNINGS**.

About Amgen

Amgen discovers, develops, manufactures and delivers innovative medicines to fight some of the world's toughest diseases. Harnessing the best of biology and technology, Amgen reaches millions of patients with its medicines.

More than 45 years ago, Amgen helped establish the biotechnology industry at its U.S. headquarters in Thousand Oaks, California, and it remains at the cutting edge of innovation, using technology and human genetic data to push beyond what is known today. Amgen is advancing a broad and deep pipeline and portfolio of medicines to treat cancer, inflammatory conditions, rare diseases, heart disease and obesity and obesity-related conditions.

Amgen has been [consistently recognized](#) for innovation and workplace culture, including honors from Fast Company and Forbes. Amgen is one of the 30 companies that comprise the Dow Jones Industrial Average®, and it is also part of the Nasdaq-100 Index®, which includes the largest and most innovative non-financial companies listed on the Nasdaq Stock Market based on market capitalization.

For more information, visit [Amgen.com](#) and follow Amgen on [X](#), [LinkedIn](#), [Instagram](#), [YouTube](#), [Facebook](#), [TikTok](#) and [Threads](#).

Amgen Forward-Looking Statements

This news release contains forward-looking statements that are based on the current expectations and beliefs of Amgen. All statements, other than statements of historical fact, are statements that could be deemed forward-looking statements, including any statements on the outcome, benefits and synergies of collaborations, or potential collaborations, with any other company (including BeOne Medicines Ltd.), the performance of Otezla® (apremilast), our acquisitions of ChemoCentryx, Inc., Dark Blue Therapeutics, Ltd. or Horizon Therapeutics plc (including the prospective performance and outlook of Horizon's business, performance and opportunities, and any potential strategic benefits, synergies or opportunities expected as a result of such acquisition), as well as estimates of revenues, operating margins, capital expenditures, cash, other financial metrics, expected legal, arbitration, political, regulatory or clinical results or practices, customer and prescriber patterns or practices, reimbursement activities and outcomes, effects of pandemics or other widespread health problems on our business, outcomes, progress, and other such estimates and results. Forward-looking statements involve significant risks and uncertainties, including those discussed below and more fully described in the Securities and Exchange Commission reports filed by Amgen, including our most recent annual report on Form 10-K and any subsequent periodic reports on Form 10-Q and current reports on Form 8-K. Unless otherwise noted, Amgen is providing this information as of the date of this news release and does not undertake any obligation to update any forward-looking statements contained in this document as a result of new information, future events or otherwise.

No forward-looking statement can be guaranteed and actual results may differ materially from those we project. Discovery or identification of new

product candidates or development of new indications for existing products cannot be guaranteed and movement from concept to product is uncertain; consequently, there can be no guarantee that any particular product candidate or development of a new indication for an existing product will be successful and become a commercial product. Further, preclinical results do not guarantee safe and effective performance of product candidates in humans. The complexity of the human body cannot be perfectly, or sometimes, even adequately modeled by computer or cell culture systems or animal models. The length of time that it takes for us to complete clinical trials and obtain regulatory approval for product marketing has in the past varied and we expect similar variability in the future. Even when clinical trials are successful, regulatory authorities may question the sufficiency for approval of the trial endpoints we have selected. We develop product candidates internally and through licensing collaborations, partnerships and joint ventures. Product candidates that are derived from relationships may be subject to disputes between the parties or may prove to be not as effective or as safe as we may have believed at the time of entering into such relationship. Also, we or others could identify safety, side effects or manufacturing problems with our products, including our devices, after they are on the market.

Our results may be affected by our ability to successfully market both new and existing products domestically and internationally, clinical and regulatory developments involving current and future products, sales growth of recently launched products, competition from other products including biosimilars, difficulties or delays in manufacturing our products and global economic conditions, including those resulting from geopolitical relations and government actions. In addition, sales of our products are affected by pricing pressure, political and public scrutiny and reimbursement policies imposed by third-party payers, including governments, private insurance plans and managed care providers and may be affected by regulatory, clinical and guideline developments and domestic and international trends toward managed care and healthcare cost containment. Furthermore, our research, testing, pricing, marketing and other operations are subject to extensive regulation by domestic and foreign government regulatory authorities. Our business may be impacted by government investigations, litigation and product liability claims. In addition, our business may be impacted by the adoption of new tax legislation or exposure to additional tax liabilities. Further, while we routinely obtain patents for our products and technology, the protection offered by our patents and patent applications may be challenged, invalidated or circumvented by our competitors, or we may fail to prevail in present and future intellectual property litigation. We perform a substantial amount of our commercial manufacturing activities at a few key facilities, including in Puerto Rico, and also depend on third parties for a portion of our manufacturing activities, and limits on supply may constrain sales of certain of our current products and product candidate development. An outbreak of disease or similar public health threat, and the public and governmental effort to mitigate against the spread of such disease, could have a significant adverse effect on the supply of materials for our manufacturing activities, the distribution of our products, the commercialization of our product candidates, and our clinical trial operations, and any such events may have a material adverse effect on our product development, product sales, business and results of operations. We rely on collaborations with third parties for the development of some of our product candidates and for the commercialization and sales of some of our commercial products. In addition, we compete with other companies with respect to many of our marketed products as well as for the discovery and development of new products. Further, some raw materials, medical devices and component parts for our products are supplied by sole third-party suppliers. Certain of our distributors, customers and payers have substantial purchasing leverage in their dealings with us. The discovery of significant problems with a product similar to one of our products that implicate an entire class of products could have a material adverse effect on sales of the affected products and on our business and results of operations. Our efforts to collaborate with or acquire other companies, products or technology, and to integrate the operations of companies or to support the products or technology we have acquired, may not be successful, and may result in unanticipated costs, delays or failures to realize the benefits of the transactions. A breakdown, cyberattack or information security breach of our information technology systems could compromise the confidentiality, integrity and availability of our systems and our data. Our stock price is volatile and may be affected by a number of events. Our business and operations may be negatively affected by the failure, or perceived failure, of achieving our sustainability objectives. The effects of global climate change and related natural disasters could negatively affect our business and operations. Global economic conditions may magnify certain risks that affect our business. Our business performance could affect or limit the ability of our Board of Directors to declare a dividend or our ability to pay a dividend or repurchase our common stock. We may not be able to access the capital and credit markets on terms that are favorable to us, or at all.

Further, any scientific information discussed in this news release relating to new indications for our products is preliminary and investigative and is not part of the labeling approved by the U.S. Food and Drug Administration for the products. The products are not approved for the investigational use(s) discussed in this news release, and no conclusions can or should be drawn regarding the safety or effectiveness of the products for these uses.

CONTACT: Amgen, Thousand Oaks
Brianna Wilkins, 513-284-9807 (media)
Casey Capparelli, 805-447-1746 (investors)

REFERENCES

1. International Agency for Research on Cancer. Trachea, bronchus and lung fact sheet. Global Cancer Observatory: Cancer Today. Published 2024. Accessed September 2, 2026. <https://gco.iarc.who.int/today/en/fact-sheets-cancers/15/trachea-bronchus-and-lung>.
2. Oronsky B, Abrouk N, Caroen S, et al. A 2022 update on extensive stage small-cell lung cancer (SCLC). *J Cancer*. 2022;13(9):2945-2953. doi:10.7150/jca.75622.
3. Sabari JK, Lok BH, Laird JH, Poirier JT, Rudin CM. Unravelling the biology of SCLC: implications for therapy. *Nat Rev Clin Oncol*. 2017;14(9):549-561. doi:10.1038/nrclinonc.2017.71.
4. Cramer-van der Welle CM, Schramel FMNH, van Leeuwen AS, Groen HJM, van de Garde EMW; Santeon SCLC Study Group. Real-world treatment patterns and outcomes of patients with extensive disease small cell lung cancer. *Eur J Cancer Care (Engl)*. 2020;29(5):e13250. doi:10.1111/ecc.13250.
5. Shaw J, Pundole X, Balasubramanian A, et al. Recent treatment patterns and real-world survival following first-line anti-PD-L1 treatment for extensive-stage small cell lung cancer. *Oncologist*. 2024;29(12):1079-1089. doi:10.1093/oncolo/loayae234.
6. Paz-Ares L, Borghaei H, Liu SV, et al. Efficacy and safety of first-line maintenance therapy with lurbinectedin plus atezolizumab in extensive-stage small-cell lung cancer (IMforte): a randomised, multicentre, open-label, phase 3 trial. *Lancet*. 2025;405(10495):2129-2143. doi:10.1016/S0140-6736(25)01011-6.
7. Reck M, Mok TSK, Mansfield A, et al. Brief report: exploratory analysis of maintenance therapy in patients with

extensive-stage SCLC treated first line with atezolizumab plus carboplatin and etoposide. *J Thorac Oncol*. 2022;17(9):1122-1129. doi:10.1016/j.jtho.2022.05.016.

8. ClinicalTrials.gov. DeLLphi-305 clinical trial listing. Accessed July 16, 2026. <https://clinicaltrials.gov/study/NCT06211036>.
9. ClinicalTrials.gov. Tarlatamab clinical trial listings. US National Library of Medicine. Accessed July 2026. <https://www.clinicaltrials.gov>.
10. Giffin MJ, Cooke K, Lobenhofer EK, et al. AMG 757, a half-life extended, DLL3-targeted bispecific T-cell engager, shows high potency and sensitivity in preclinical models of small-cell lung cancer. *Clin Cancer Res*. 2021;27(5):1526-1537. doi:10.1158/1078-0432.CCR-20-2845.
11. Baeuerle PA, Kufer P, Bargou R. BiTE: Teaching antibodies to engage T-cells for cancer therapy. *Curr Opin Mol Ther*. 2009;11(1):22-30.
12. Ahn MJ, Cho BC, Felip E, et al. Tarlatamab for patients with previously treated small-cell lung cancer. *N Engl J Med*. 2023;389(22):2063-2075. doi:10.1056/NEJMoa2307980/
13. Rojo F, Corassa M, Mavroudis D, et al. International real-world study of DLL3 expression in patients with small cell lung cancer. *Lung Cancer*. 2020;147:237-243. doi:10.1016/j.lungcan.2020.07.026.



 View original content to download multimedia: <https://www.prnewswire.com/news-releases/indelltra-in-combination-with-imfinzi-demonstrated-landmark-improvement-in-overall-survival-in-first-line-extensive-stage-small-cell-lung-cancer-302872305.html>

SOURCE Amgen